


What is claimed is:

1. A compound of the formula I



or a stereoisomeric form or a pharmaceutically acceptable salt of the compound of the formula I, wherein

A is $-(C_1-C_6)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

- O- R^1 or
- C(O)-OR¹,
- C(O)-NR¹R¹,
- C(O)-NR¹-SO₂R¹,
- NR¹R¹,
- CN, in which R^1 is

hydrogen,

$-(C_1-C_6)$ -alkyl,

$-(C_6-C_{14})$ aryl or

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,

-O- R^1 ,

-SR¹,

-S(O)- R^1

-S(O)₂- R^1

-C(O)-OR¹,

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is $2n + 1$,

5

$-C(O)-NR^1R^1$,
 $-C(O)-NR^1-SO_2R^1$,
 $-NR^1R^1$,
 $-CN$,

10

heteroaryl having from 5 to 14 ring members, in which heteroaryl is unsubstituted or optionally substituted once or more, independently of each other, by R^2 , in which R^2 is

15

$-(C_1-C_4)$ -alkyl,
 $-OH$,
 $-O-(C_1-C_4)$ -alkyl,
halogen,
 $-N(R^3)-R^4$, in which R^3 and R^4 are, independently of each other, hydrogen atom or $-(C_1-C_4)$ -alkyl,

20

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

25


$-CN$,
 $-SR^1$,
 $-S(O)-R^1$,
 $-S(O)_2-R^1$ or
 $-C(O)-NR^1R^1$,

30

$-(C_3-C_6)$ -cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

B is a covalent bond,



$-\text{O}(\text{CH}_2)_a-$, in which a is an integer from 1 to 4,

O, S, NR^2 , $-\text{C}(\text{O})-$, $-\text{NR}^2-\text{C}(\text{O})-$, $-\text{C}(\text{O})-\text{NR}^2-$, $-\text{NR}^2-\text{SO}_2-$,
 5 $-\text{SO}_2-\text{NR}^2-$, $-\text{NR}^2-\text{C}(\text{O})-\text{NR}^2-$, and R^2 is defined as above, or

$-(\text{C}_1-\text{C}_4)\text{-alkylene}$, in which alkylene is straight-chain or
 branched and is optionally substituted, once or more,
 independently of each other, by R^1 , and R^1 is defined as above,

10 D is $-(\text{C}_1-\text{C}_6)\text{-alkyl}$, in which alkyl is straight-chain or branched and is
 optionally substituted, once or more, independently of each
 other, by R^1 , and R^1 is defined as above,

15 heteroaryl having from 5 to 14 ring members, in which heteroaryl
 is unsubstituted or is substituted once or more, independently of
 each other, by R^2 , and R^2 is defined as above,

20 heterocycle having from 5 to 12 ring members, in which
 heterocycle is unsubstituted or substituted, once or more,
 independently of each other, by R^2 , and R^2 is defined as above,

25 $-(\text{C}_6-\text{C}_{14})\text{-aryl}$, in which aryl is unsubstituted or substituted, once
 or more, independently of each other, by R^2 , and R^2 is defined
 as above, or

$-(\text{C}_3-\text{C}_6)\text{-cycloalkyl}$, in which cycloalkyl is unsubstituted or
 substituted, once or more, independently of each other, by R^2 ,
 and R^2 is defined as above, or

30 B-D is hydrogen,
 halogen,

fluoroalkyl of the formula $-\text{C}_n\text{H}_x\text{F}_y$ or fluoroalkoxy of the formula
 $-\text{OC}_n\text{H}_x\text{F}_y$, wherein n is an integer from 1 to 4, x is an integer
 from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n +$
 1,

-(CH₂)_a-Y-R³, in which a is an integer from 1 to 4, Y is O, S, NR², and R³ is

-(C₁-C₆)-alkyl,

-(C₆-C₁₄)-aryl,

5 -(C₃-C₆)-cycloalkyl, and

R is hydrogen,

-(C₁-C₆)-alkyl, or

10 -(C₆-C₁₄)-aryl-(C₁-C₆)-alkyl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, and

X and Z are identical or different and are, independently of each other selected from:

15 hydrogen atom,

-(C₁-C₄)-alkyl,

-OH,

-O-(C₁-C₄-alkyl),

halogen,

20 fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is 2n + 1,

-C(O)-OR¹,

25 -C(O)-NR¹R¹,

-C(O)-NR¹-SO₂R¹,

-NR¹R¹,

-NR¹-C(O)-NR¹R¹,

-NR¹-C(O)-R¹,

30 -NR¹-C(O)-OR¹,

-O-C(O)-NR¹R¹,

-CN,

-SR¹,

-S(O)-R¹,

35 -S(O)₂-R¹,

-S(O)₂-NR¹R¹,

-NR¹-SO₂-R¹, in which R¹ is as defined above,

heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

with the proviso that when A is -(C₁-C₆)-alkyl, -O-R¹, -C(O)-OR¹, or heteroaryl, at least one of the following applies:

B is not a covalent bond or -(C₁-C₄)-alkylene,

D is not heteroaryl, heterocycle, -(C₆-C₁₄)-aryl, -(C₃-C₆)-cycloalkyl, or

X and Z are not -(C₁-C₄)-alkyl, -OH, -O-(C₁-C₄)-alkyl, or halogen.

2. A compound of the formula I as claimed in claim 1, wherein

A is -(C₁-C₃)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

-O-R¹, or

-C(O)-OR¹, in which R¹ is

hydrogen,

-(C₁-C₃)-alkyl, or

-CF₃

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 3, x is an integer from 0 to 6, y is an integer from 1 to 7 and sum of x and y is 2n + 1,

B is a covalent bond or O,

D is phenyl or naphthyl, in which phenyl or naphthyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , in which R^2 is

fluorine, chlorine or bromine,

-OH,

-CF₃,

-SR¹, in which R¹ is defined as above,

-(C₁-C₄)-alkyl

-O-(C₁-C₂)-alkyl or

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₃)-alkyl,

heteroaryl selected from the group consisting of pyridyl, furanyl, pyrrolyl, isoxazolyl, benzofuranyl, benzothiophenyl, quinolinyl, isoquinolinyl, quinoxalinyl and thiophenyl, in which heteroaryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , in which R^2 is defined as above or

-(C₄-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

B-D is ((CH₂)_a-Y-R³, in which a is an integer from 1 to 2, Y is O and R³ is -(C₁-C₃)-alkyl, and

R is hydrogen,

-(C₁-C₃)-alkyl, or

-phenyl-(C₁-C₃)-alkyl, and

X and Z are identical or different and are, independently of each other, hydrogen, -C(O)-O(C₁-C₃)alkyl, -OCH₃, -N(CH₃)₂ or halogen.

3. A compound of the formula 1 as claimed in claim 1, wherein the compound of the formula I is selected from the group consisting of:
- 5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
 - 3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,

1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-
isoquinoline,
5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-
5 isoquinoline,
1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
10 7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
15 7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
6-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
20 8-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-
isoquinoline,
25 3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-
isoquinoline, and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-
isoquinoline.

30


4. A compound selected from the group consisting of:
5-benzo[b]thiophene-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
35 5-(3-chloro-benzo[b]thiophen-2-yl)-3-methyl-1H-pyrazolo[4,3-c]-
isoquinoline,

- 5-(2-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
5-(6-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
5 3-ethyl-5-pyridin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
5-furan-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-methoxy-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-benzofuran-2-yl)-1H-pyrazolo[4,3-c]-
isoquinoline,
10 3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
15 9-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
20 3-methyl-5-(4-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
25 3-methyl-5-(2,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
30 3-methyl-5-(2,4-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
6-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
35 8-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,

- 3,9-dimethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(1-methyl-1H-pyrrol-2-yl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
5 3-methyl-5-quinoxalin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-thiophen-2-yl-1H-pyrazolo[4,3-c]isoquinoline, and
3-methyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]isoquinoline.
5. A pharmaceutical composition comprising a therapeutically effective
10 content of at least one compound of the formula I as claimed in
claim 1 together with a pharmaceutically suitable carrier optionally in
combination with a suitable additive, other active compounds and
auxiliary substances.
- 15 6. A method of treating a disease condition associated with the
increased activity of NIK comprising administering to a patient
suffering from said disease condition a therapeutically effective
amount of a compound according to claim 1.
- 20 7. The method as claimed in claim 6, wherein the compound is
according to claim 2.
8. The method as claimed in claim 6 wherein said compound is
selected from the group consisting of:
- 25 5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-
30 isoquinoline,
5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-
isoquinoline,
1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
35 1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,

7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
5 7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-
isoquinoline,
10 3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-
isoquinoline, and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-
isoquinoline.

- 15 9. The method as claimed in claim 6, wherein the disease condition is
caused due to an inflammatory component.
- 20 10. The method as claimed in claim 6, wherein the diseases are
osteoarthritis, rheumatoid arthritis, asthma, irritable bowel disease,
Alzheimer's disease, stroke, diabetes, atherosclerosis, multiple
sclerosis, rejection reactions on the part of the body against a
transplanted organ or rejection reactions on the part of the
transplanted organ against the body.
- 25 11. A pharmaceutical composition comprising a compound of the
formula (I)



(I)

30 or a stereoisomeric form or a pharmaceutically acceptable salt of the
compound of the formula I, wherein

A is $-(C_1-C_6)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

5 $-O-R^1$ or
 $-C(O)-OR^1$,
 $-C(O)-NR^1R^1$,
 $-C(O)-NR^1-SO_2R^1$,
 $-NR^1R^1$,

10 $-CN$, in which R^1 is

 hydrogen,

$-(C_1-C_6)$ -alkyl,

$-(C_6-C_{14})$ aryl or

 fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the
15 formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6,
 x is an integer from 0 to 12, y is an integer from 1 to 13
 and sum of x and y is $2n + 1$,

$-O-R^1$,

$-SR^1$,

20 $-S(O)-R^1$

$-S(O)_2-R^1$

$-C(O)-OR^1$,

 fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula

$-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer
25 from 0 to 12, y is an integer from 1 to 13 and sum of x and y is
 $2n + 1$,

$-C(O)-NR^1R^1$,

$-C(O)-NR^1-SO_2R^1$,

$-NR^1R^1$,

30 $-CN$,

 heteroaryl having from 5 to 14 ring members, in which heteroaryl
is unsubstituted or optionally substituted once or more,
independently of each other, by R^2 , in which R^2 is

$-(C_1-C_4)$ -alkyl,

- OH,
 -O-(C₁-C₄)-alkyl,
 halogen,
 -N(R³)-R⁴, in which R³ and R⁴ are, independently of each
 5 other, hydrogen atom or -(C₁-C₄)-alkyl,
 fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the
 formula -OC_nH_xF_y, wherein n is an integer from 1 to 4, x is an
 integer from 0 to 8, y is an integer from 1 to 9 and sum of x
 and y is 2n + 1,
 10 -CN,
 -SR¹,
 -S(O)-R¹,
 -S(O)₂-R¹ or
 -C(O)-NR¹R¹,
 15 -(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or
 substituted, once or more, independently of each other, by R²,
 and R² is defined as above,
 heterocycle having from 5 to 12 ring members, in which
 heterocycle is unsubstituted or substituted, once or more,
 20 independently of each other, by R², and R² is defined as above,
 B is a covalent bond,
 -C=CR¹-,
 —C≡C—,
 25 -O(CH₂)_a-, in which a is an integer from 1 to 4,
 O, S, NR², -C(O)-, -NR²-C(O)-, -C(O)-NR²-, -NR²-SO₂-,
 -SO₂-NR²-, -NR²-C(O)-NR²-, and R² is defined as above, or
 -(C₁-C₄)-alkylene, in which alkylene is straight-chain or
 branched and is optionally substituted, once or more,
 30 independently of each other, by R¹, and R¹ is defined as above,
 D is -(C₁-C₆)-alkyl, in which alkyl is straight-chain or branched and is
 optionally substituted, once or more, independently of each
 other, by R¹, and R¹ is defined as above,
 35

heteroaryl having from 5 to 14 ring members, in which heteroaryl is unsubstituted or is substituted once or more, independently of each other, by R^2 , and R^2 is defined as above,

5 heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

10 $-(C_6-C_{14})$ -aryl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

15 $-(C_3-C_6)$ -cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

B-D is hydrogen,
halogen,

20 fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

$-(CH_2)_a-Y-R^3$, in which a is an integer from 1 to 4, Y is O, S, NR^2 , and R^3 is

25 $-(C_1-C_6)$ -alkyl,
 $-(C_6-C_{14})$ -aryl,
 $-(C_3-C_6)$ -cycloalkyl, and

R is hydrogen,

30 $-(C_1-C_6)$ -alkyl, or
 $-(C_6-C_{14})$ -aryl- (C_1-C_6) -alkyl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, and

X and Z are identical or different and are, independently of each other selected from:

- hydrogen atom,
- (C₁-C₄)-alkyl,
- 5 -OH,
- O-(C₁-C₄-alkyl),
- halogen,
- fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer
- 10 from 0 to 12, y is an integer from 1 to 13 and sum of x and y is 2n + 1,
- C(O)-OR¹,
- C(O)-NR¹R¹,
- C(O)-NR¹-SO₂R¹,
- 15 -NR¹R¹,
- NR¹-C(O)-NR¹R¹,
- NR¹-C(O)-R¹,
- NR¹-C(O)-OR¹,
- O-C(O)-NR¹R¹,
- 20 -CN,
- SR¹,
- S(O)-R¹,
- S(O)₂-R¹,
- S(O)₂-NR¹R¹,
- 25 -NR¹-SO₂-R¹, in which R¹ is as defined above,
- heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or
- 30 -(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

with the proviso that when A is -(C₁-C₆)-alkyl, -O-R¹, -C(O)-OR¹, or

heteroaryl, at least one of the following applies:

B is not a covalent bond or $-(C_1-C_4)$ -alkylene,

D is not heteroaryl, heterocycle, $-(C_6-C_{14})$ -aryl, $-(C_3-C_6)$ -cycloalkyl, or

5 X and Z are not $-(C_1-C_4)$ -alkyl, -OH, $-O-(C_1-C_4)$ -alkyl, or halogen.

12. The composition as claimed in claim 11, wherein

10 A is $-(C_1-C_3)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

$-O-R^1$, or

$-C(O)-OR^1$, in which R^1 is

hydrogen,

15 $-(C_1-C_3)$ -alkyl, or

$-CF_3$

20 fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 3, x is an integer from 0 to 6, y is an integer from 1 to 7 and sum of x and y is $2n + 1$,

B is a covalent bond or O,

25 D is phenyl or naphthyl, in which phenyl or naphthyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , in which R^2 is

fluorine, chlorine or bromine,

-OH,

$-CF_3$,

30 $-SR^1$, in which R^1 is defined as above,

$-(C_1-C_4)$ -alkyl

$-O-(C_1-C_2)$ -alkyl or

$-N(R^3)-R^4$, in which R^3 and R^4 are, independently of each other, hydrogen atom or $-(C_1-C_3)$ -alkyl,

heteroaryl selected from the group consisting of pyridyl, furanyl, pyrrolyl, isoxazolyl, benzofuranyl, benzothiophenyl, quinoliny, isoquinoliny, quinoxaliny and thiophenyl, in which heteroaryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , in which R^2 is defined as above or

-(C₄-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

B-D is ((CH₂)_a-Y-R³, in which a is an integer from 1 to 2, Y is O and R³ is -(C₁-C₃)-alkyl, and

R is hydrogen,

-(C₁-C₃)-alkyl, or

-phenyl-(C₁-C₃)-alkyl, and

X and Z are identical or different and are, independently of each other, hydrogen, -C(O)-O(C₁-C₃)alkyl, -OCH₃, -N(CH₃)₂ or halogen.

13. The composition as claimed in claim 11, wherein the compound of the formula I is selected from the group consisting of:

5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,

3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,

3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,

3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,

1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,

5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,

1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-isoquinoline,

1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,

1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,


1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,

5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,

7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-isoquinoline,

7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
5 isoquinoline,
6-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
8-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
10 8-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-
isoquinoline,
3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
15 3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-
isoquinoline, and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-
isoquinoline.

- 20 14. A method of treating a disease condition associated with
inflammation comprising administering to a patient suffering from
said disease condition a therapeutically effective amount of a
compound of formula (I):



(I)

25 or a stereoisomeric form or a pharmaceutically acceptable salt of the
compound of the formula I, wherein

A is $-(C_1-C_6)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

$-O-R^1$ or
 $-C(O)-OR^1$,
 $-C(O)-NR^1R^1$,
 $-C(O)-NR^1-SO_2R^1$,
 $-NR^1R^1$,
 $-CN$, in which R^1 is

hydrogen,
 $-(C_1-C_6)$ -alkyl,
 $-(C_6-C_{14})$ aryl or
fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the
formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 6,
 x is an integer from 0 to 12, y is an integer from 1 to 13
and sum of x and y is $2n + 1$,

$-O-R^1$,
 $-SR^1$,
 $-S(O)-R^1$,
 $-S(O)_2-R^1$,
 $-C(O)-OR^1$,

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula
 $-OC_nH_xF_y$, wherein n is an integer from 1 to 6, x is an integer
from 0 to 12, y is an integer from 1 to 13 and sum of x and y is
 $2n + 1$,

$-C(O)-NR^1R^1$,
 $-C(O)-NR^1-SO_2R^1$,
 $-NR^1R^1$,
 $-CN$,

heteroaryl having from 5 to 14 ring members, in which heteroaryl
is unsubstituted or optionally substituted once or more,
independently of each other, by R^2 , in which R^2 is

$-(C_1-C_4)$ -alkyl,
 $-OH$,

-O-(C₁-C₄)-alkyl,
halogen,

-N(R³)-R⁴, in which R³ and R⁴ are, independently of each other, hydrogen atom or -(C₁-C₄)-alkyl,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula -OC_nH_xF_y, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is 2n + 1,

-CN,
-SR¹,

-S(O)-R¹,

-S(O)₂-R¹ or

-C(O)-NR¹R¹,

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above,

B is a covalent bond,

-C=CR¹-,

—C≡C—

-O(CH₂)_a-, in which a is an integer from 1 to 4,

O, S, NR², -C(O)-, -NR²-C(O)-, -C(O)-NR²-, -NR²-SO₂-, -SO₂-NR²-, -NR²-C(O)-NR²-, and R² is defined as above, or

-(C₁-C₄)-alkylene, in which alkylene is straight-chain or branched and is optionally substituted, once or more, independently of each other, by R¹, and R¹ is defined as above,

D is -(C₁-C₆)-alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by R¹, and R¹ is defined as above,

heteroaryl having from 5 to 14 ring members, in which heteroaryl is unsubstituted or is substituted once or more, independently of each other, by R^2 , and R^2 is defined as above,

5 heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above,

10 $-(C_6-C_{14})$ -aryl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

15 $-(C_3-C_6)$ -cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, or

B-D is hydrogen,
halogen,

20 fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 4, x is an integer from 0 to 8, y is an integer from 1 to 9 and sum of x and y is $2n + 1$,

$-(CH_2)_a-Y-R^3$, in which a is an integer from 1 to 4, Y is O, S, NR^2 , and R^3 is

25 $-(C_1-C_6)$ -alkyl,
 $-(C_6-C_{14})$ -aryl,
 $-(C_3-C_6)$ -cycloalkyl, and

R is hydrogen,

30 $-(C_1-C_6)$ -alkyl, or
 $-(C_6-C_{14})$ -aryl- $-(C_1-C_6)$ -alkyl, in which aryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , and R^2 is defined as above, and

X and Z are identical or different and are, independently of each other selected from:

hydrogen atom,

-(C₁-C₄)-alkyl,

5

-OH,

-O-(C₁-C₄-alkyl),

halogen,

fluoroalkyl of the formula -C_nH_xF_y or fluoroalkoxy of the formula

10

-OC_nH_xF_y, wherein n is an integer from 1 to 6, x is an integer from 0 to 12, y is an integer from 1 to 13 and sum of x and y is 2n + 1,

-C(O)-OR¹,

-C(O)-NR¹R¹,

15

-C(O)-NR¹-SO₂R¹,

-NR¹R¹,

-NR¹-C(O)-NR¹R¹,

-NR¹-C(O)-R¹,

-NR¹-C(O)-OR¹,

-O-C(O)-NR¹R¹,

20

-CN,

-SR¹,

-S(O)-R¹,

-S(O)₂-R¹,

-S(O)₂-NR¹R¹,

25

-NR¹-SO₂-R¹, in which R¹ is as defined above,

heterocycle having from 5 to 12 ring members, in which heterocycle is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

30

-(C₃-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above.

15. The method as claimed in claim 14 wherein

A is $-(C_1-C_3)$ -alkyl, in which alkyl is straight-chain or branched and is optionally substituted, once or more, independently of each other, by

$-O-R^1$, or

$-C(O)-OR^1$, in which R^1 is

hydrogen,

$-(C_1-C_3)$ -alkyl, or

$-CF_3$

fluoroalkyl of the formula $-C_nH_xF_y$ or fluoroalkoxy of the formula $-OC_nH_xF_y$, wherein n is an integer from 1 to 3, x is an integer from 0 to 6, y is an integer from 1 to 7 and sum of x and y is $2n + 1$,

B is a covalent bond or O,

D is phenyl or naphthyl, in which phenyl or naphthyl is unsubstituted or substituted, once or more, independently of each other, by R^2 , in which R^2 is

fluorine, chlorine or bromine,

$-OH$,

$-CF_3$,

$-SR^1$, in which R^1 is defined as above,

$-(C_1-C_4)$ -alkyl

$-O-(C_1-C_2)$ -alkyl or

$-N(R^3)-R^4$, in which R^3 and R^4 are, independently of each other, hydrogen atom or $-(C_1-C_3)$ -alkyl,

heteroaryl selected from the group consisting of pyridyl, furanyl, pyrrolyl, isoxazolyl, benzofuranyl, benzothiophenyl, quinolinyl, isoquinolinyl, quinoxalinyl and thiophenyl, in which heteroaryl is unsubstituted or substituted, once or more, independently of each other, by R^2 , in which R^2 is defined as above or

-(C₄-C₆)-cycloalkyl, in which cycloalkyl is unsubstituted or substituted, once or more, independently of each other, by R², and R² is defined as above, or

B-D is ((CH₂)_a-Y-R³), in which a is an integer from 1 to 2, Y is O and R³ is -(C₁-C₃)-alkyl, and

R is hydrogen,

-(C₁-C₃)-alkyl, or

-phenyl-(C₁-C₃)-alkyl, and

X and Z are identical or different and are, independently of each other, hydrogen, -C(O)-O(C₁-C₃)alkyl, -OCH₃, -N(CH₃)₂ or halogen.

16. The method as claimed in claim 14, wherein the compound of formula (I) is selected from the group consisting of:

5-pyridin-2-yl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-trifluoromethyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-
isoquinoline,
5-phenyl-3-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
1,3-dimethyl-5-(3-trifluoromethylphenyl)-1H-pyrazolo[4,3-c]-
isoquinoline,
1,3-dimethyl-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-cyclohexyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
1-benzyl-5-naphthyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
5-methoxymethyl-3-methyl-1H-pyrazolo[4,3-c]-isoquinoline,
7-methoxycarbonyl-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-methoxycarbonyl-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
7-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
6-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,

- 6-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
isoquinoline,
8-dimethylamino-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-dimethylamino-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]-
5 isoquinoline,
1,3-dimethyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]-
isoquinoline,
3-methyl-5-phenyl-9-trifluoromethyl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-pyridin-2-yl-9-trifluoromethyl-1H-pyrazolo[4,3-c]-
10 isoquinoline, and
3-methyl-5-(2,3,4,5,6-pentafluoro-phenyl)-1H-pyrazolo[4,3-c]-
isoquinoline.
17. The method as claimed in claim 14, wherein the compound of
15 formula (I) is selected from the group consisting of:
5-benzo[b]thiophene-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
7-bromo-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
5-(3-chloro-benzo[b]thiophen-2-yl)-3-methyl-1H-pyrazolo[4,3-c]-
20 isoquinoline,
5-(2-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
5-(6-chloro-pyridin-3-yl)-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3-ethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
25 3-ethyl-5-pyridin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
5-furan-2-yl-3-methyl-1H-pyrazolo[4,3-c]isoquinoline,
7-methoxy-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-benzofuran-2-yl)-1H-pyrazolo[4,3-c]-
isoquinoline,
30 3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
35 9-chloro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
9-fluoro-3-methyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-phenyl-1H-pyrazolo[4,3-c]isoquinoline,

- 3-methyl-5-(2-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-methyl-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
5 3-methyl-5-(3-bromo-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-chloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(3,4-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
10 3-methyl-5-(2,6-dichloro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(4-fluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,4-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(2,6-difluoro-phenyl)-1H-pyrazolo[4,3-c]isoquinoline,
15 6-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
6-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
8-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
9-chloro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
20 9-fluoro-3-methyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3,9-dimethyl-5-pyridin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-(1-methyl-1H-pyrrol-2-yl)-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-quinolin-3-yl-1H-pyrazolo[4,3-c]isoquinoline,
25 3-methyl-5-quinoxalin-2-yl-1H-pyrazolo[4,3-c]isoquinoline,
3-methyl-5-thiophen-2-yl-1H-pyrazolo[4,3-c]isoquinoline, and
3-methyl-5-(3-methyl-thiophen-2-yl)-1H-pyrazolo[4,3-c]isoquinoline.
18. The method as claimed in claim 14, wherein the disease condition is
30 selected from the group consisting of multiple sclerosis,
atherosclerosis, inflammatory bowel disease, Alzheimer's disease,
stroke and diabetes.
19. The method as claimed in claim 18, wherein the disease condition is
35 multiple sclerosis.

20. The method as claimed in claim 18, wherein the disease condition is atherosclerosis.
21. The method as claimed in claim 18, wherein the disease condition is inflammatory bowel disease.
22. The method as claimed in claim 18, wherein the disease condition is Alzheimer's disease.
23. The method as claimed in claim 18, wherein the disease condition is stroke.
24. The method as claimed in claim 18, wherein the disease condition is diabetes.